

**SUMMARY OF LANDSCAPE-MANAGEMENT COMPARTMENTS, PAST AND CURRENT MANAGEMENT AND OBJECTIVES.**

Area name	Numbered compartments	Habitat type	Wildlife features	Conservation rating	Previous Management (2005-2009)	Current management
<b>Ceremonial Way</b>	<b>1,2,3</b>	Unimproved grassland	glaucous sedge, field scabious, bird's-foot trefoil, lady's bedstraw and creeping restharrow	High	Initial: large-scale tree-felling (ash and sycamore), stump-treatment.	Regular grass-cutting (Rest), regular cutting of grave areas and vegetation; sapling-thinning (ash and sycamore) and stump
<b>Bath Road Frontage</b>	<b>4,5</b>	Scrub		Low	Initial: large-scale scrub clearance	Rotational management over 10 years
		Secondary woodland	English elm, Ivy broomrape	Medium	Initial: improvements to canopy structure; tree-felling and stump treatment,	Footpath maintenance; sapling -thinning (ash and sycamore) and stump
<b>Poplar Ridge</b>	<b>6,7,8</b>	Species-rich grassland	Upright brome, field scabious, glaucous sedge, lady's bedstraw, creeping restharrow, rough hawkbit. Several nationally rare and scarce species of invertebrate Slow worms	High	Initial: large-scale tree-felling (ash and sycamore), stump-treatment.	Annual meadow cutting; scrub-encroachment control; sapling -thinning (ash and sycamore), stump treatment
		Scrub	Wild privet, field rose, breeding birds (blackcap)	Low	Initial: large-scale scrub clearance	Rotational management over 10 years
		Secondary woodland	White poplar (standing and deadwood); eight species of nationally rare or scarce invertebrate; ivy broomrape, breeding birds	High	Initial: improvements to canopy structure; tree-felling (ash and sycamore), stump-treatment. Japanese Knotweed eradication.	Footpath maintenance; sapling -thinning (ash and sycamore) and stump treatment of Japanese Knotweed; re-growth.
		Species-diverse grassland fringe	Grey sedge, primrose, crow garlic, germander speedwell.	Medium	Initial: scrub clearance	Annual meadow cutting; scrub-encroachment control; sapling -thinning (ash and sycamore), stump treatment

**SUMMARY OF LANDSCAPE-MANAGEMENT COMPARTMENTS, PAST AND CURRENT MANAGEMENT AND OBJECTIVES.**

<b>Area name</b>	<b>Numbered compartments</b>	<b>Habitat type</b>	<b>Wildlife features</b>	<b>Conservation rating</b>	<b>Previous Management (2005-2009)</b>	<b>Current management</b>
<b>Combe</b>		Secondary woodland	Ramsons, breeding birds (chiff-chaff)	Medium	Initial: improvements to canopy structure; tree-felling (ash and sycamore), stump-treatment. Japanese Knotweed eradication.	Footpath maintenance, sapling -thinning = a treatment, treatment. Knotweed re-growth
<b>Upper Combe Slopes</b>	<b>14,15,16</b>	Grassland		Low	Annual meadow cut	Annual meadow cut
		Secondary woodland	wild valerian, goldilocks buttercup, ivy broomrape.	Medium	Initial: improvements to canopy structure; tree-felling (ash and sycamore), stump-treatment.	footpath maintenance, sapling -thinning (ash and sycamore) and re-rotation year rotation
		Scrub	wild privet, apple, field maple, Ivy broomrape, Breeding birds (blackcap, long-tailed tit).	Medium	Initial: large-scale scrub clearance	Rotational management every 5 years
<b>Lower Combe Slopes</b>	<b>17,18</b>	Species-rich grassland	field scabious, creeping restharrow, glaucous sedge, meadow vetchling, spiny restharrow, zig-zag clover, birdsfoot trefoil, burnet saxifrage	High	Annual meadow cut and footpath maintenance	Annual meadow cut and scrub-encroachment, sapling–thinning, st
		Secondary woodland	Badger sett, horse chestnut, various conifers, field maple, wild privet , holly	Medium	Initial: improvements to canopy structure; tree-felling (ash and sycamore), stump-treatment.	Footpath maintenance, sapling -thinning (ash and sycamore) and stump-treatment every 5 years,
<b>Lower Slopes</b>	<b>19,20,21,22</b>	Scrub	Breeding birds (blackcap, bullfinch)	Medium	Initial: selective scrub-removal	Rotational management every 5 years
<b>Upper Slopes</b>	<b>23,24,25,26</b>	Species-rich grassland fringe	field scabious, creeping restharrow, bird’s-foot trefoil, lady’s bedstraw and zig-zag	High	Annual meadow cut	Annual meadow cut and scrub-encroachment, sapling -thinning (ash

**SUMMARY OF LANDSCAPE-MANAGEMENT COMPARTMENTS, PAST AND CURRENT MANAGEMENT AND OBJECTIVES.**

Area name	Numbered compartments	Habitat type	Wildlife features	Conservation rating	Previous Management (2005-2009)	Current management
			primrose, Ivy broomrape, breeding birds (jay, stock dove, chiffchaff, sparrowhawk)			rotation, Japanese knotweed eradication.
		Species-rich grassland patch in Comp 26	Common knapweed, cuckoo flower, soapwort	High	Annual meadow cut	Annual meadow cut
<b>Upper Cemetery</b>	<b>27,28,29,30</b>	Secondary woodland		Low	Initial: Japanese Knotweed eradication.	Thinning of ash and saplings and trees, stump treatment, footpath
		Grassland	Slow worms, lesser stitchwort, meadow vetchling, nationally rare and scarce species of invertebrate ( <i>Momphus bradleyi</i> , and <i>Tricimba humeralis</i> ).	Medium	<b>Area outside of HLF-funded zone – no other initial restoration works carried out.</b>	Annual meadow cut (year), regular footpath maintenance, grass-ramble-cutting, selective thinning, stump treatment
		Scrub	Breeding birds (long-tailed tit, bullfinch, blackcap, lesser whitethroat)	Medium		Rotational 'scallop' edge in selected areas
<b>Old Orchard</b>	<b>31</b>	Secondary woodland	semi-mature ash, sycamore, apple and pedunculate oak, hazel, gooseberry, blackthorn, ramsons, sweet violet, hairy brome, hart's-tongue fern, male fern and soft shield fern, breeding birds (jay, great-spotted woodpecker)	High	<b>Area outside of HLF-funded zone – no other initial restoration works carried out.</b>	Health and safety management
		Scrub fringe	Butterflies, scarce hoverflies ( <i>Volucella zonaria</i> and	High		Footpath maintenance